



Metal stress response influences a soil bacterial community's permissiveness towards a broad-host-range plasmid

Klümper, Uli; Brandt, Kristian Koefoed; Dechesne, Arnaud; Riber, Leise; Sørensen, Søren J.; Smets, Barth F.

Published in:
The danish microbiological society - Annual congress 2014

Publication date:
2014

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Klümper, U., Brandt, K. K., Dechesne, A., Riber, L., Sørensen, S. J., & Smets, B. F. (2014). Metal stress response influences a soil bacterial community's permissiveness towards a broad-host-range plasmid. In *The danish microbiological society - Annual congress 2014: Program & abstracts* (pp. 20-20). American Society for Microbiology.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Dominant clones of livers, spleens and fecal samples were whole genome sequenced to analyze for the presence of adaptive mutations. In one mouse a dominant clone from the fecal sample had a non-synonymous point mutation in a regulator of central carbon metabolism. In another mouse, dominant clones from both the liver and fecal sample had the same non-synonymous point mutation in a regulator of virulence gene expression. Interestingly, this clone was selected against in the spleen-sample from the same mouse. The contribution of these mutations to virulence and host-adaptation of *Salmonella* is presently being investigated. In conclusion, even during a short "chronic" infection *Salmonella* undergoes genetic adaptation to its host. Furthermore, the genetic and host-niche specific nature of the mutations indicate that adjustment of central metabolism and virulence regulation involves niche-specific selection.

P2: Metal stress response influences a soil bacterial community's permissiveness towards a broad host range plasmid

Uli Klümper¹, Kristian K. Brandt², Arnaud Dechesne¹, Leise Riber³, Søren J. Sørensen³, Barth F. Smets¹

¹ DTU Environment, Technical University of Denmark, ² Department of Plant and Environmental Sciences, University of Copenhagen, ³ Department of Biology, University of Copenhagen

The extent by which antibiotic resistance encoding plasmids transfer is of acute relevance in the age of massive antibiotic usage. The occurrence of stressors might play a major role in altering the acute permissiveness of a community towards plasmids, since plasmid transfer is considered a main process in immediate stress response and adaptation. A major stress factor for soil communities is the introduction of metals through geological or anthropogenic sources like manure.

We, therefore, aimed to assess how the introduction of metal stress alters a soil community's permissiveness towards plasmids. We also tried to evaluate if a general or a metal specific stress response exists.

Hence, using a ³H-leucine-incorporation approach, we measured 25% and 50% inhibition concentrations for 5 metals (Cu, Zn, Ni, Cd, As). A mCherry-tagged *E. coli* donor carrying the *gfp*-tagged plasmid pJK5 was mated with a soil bacterial community and exposed to metal stress. Plasmid transfer was quantified by detecting green fluorescent transconjugants. Transconjugants were isolated using fluorescent activated cell sorting. Sorted transconjugants were analyzed by 16S rRNA gene amplicon pyrosequencing. Within all transconjugal pools a high diversity of transconjugants remained irrespective of stress exposure. Still, results revealed an effect in soil microbial community permissiveness towards broad-host-range plasmids when exposed to the heavy metals Ni or Cu, while As exposure showed no effect.

**The Danish Microbiological Society
Annual Congress 2014
Program & Abstracts**

**Monday, 10th November 2014
Eigtved's Pakhus
Copenhagen**



**AMERICAN
SOCIETY FOR
MICROBIOLOGY**

